

Abstract

In 2017, areas of northern New South Wales experienced significant flooding as a result of ex-Tropical Cyclone Debbie. Such events are likely to become more frequent and severe due to climate change. There is a current gap in the literature investigating the effects of indirect disruption caused by flooding (e.g. loss of access to health and social care for people who have not had their property inundated) on mental health. A survey was conducted of flood-affected communities across northern New South Wales six months after the event to investigate relationships between the flood and adverse mental health outcomes. Responses were used to investigate associations between indirect disruptions and psychological morbidity. Respondents who reported indirect disruption were significantly more likely to report experiences consistent with probable post-traumatic stress disorder than those who did not report any disruption. Those who reported a loss of health and social care or a disruption to their utilities were more likely to experience adverse mental health outcomes. This study showed that indirect disruption due to flooding is associated significantly with adverse mental health. Post-disaster recovery managers might consider allocating mental health support for people who have experienced indirect disruptions.

Disruptions and mental-health outcomes following Cyclone Debbie

Jonathan King^{1,2}

Dr Jo Longman¹

Dr Veronica Matthews¹

Professor James Bennett-Levy¹

Professor Ross S Bailie¹

Steve Carrig³

Associate Professor Megan Passey¹

¹ University of Sydney, Lismore, New South Wales.

² University of New South Wales, Sydney, New South Wales.

³ New South Wales Health, Lismore, New South Wales.

SUBMITTED

6 August 2019

ACCEPTED

9 December 2019

Introduction

Recent years have seen a growing literature on the mental health and wellbeing of individuals affected by flooding (Fernandez *et al.* 2015). Much of the focus has been on the relationship between direct disruption due to flooding and poor mental health (Alderman, Turner & Tong 2012; Zhong *et al.* 2018; Waite *et al.* 2017; Jermacane *et al.* 2018; Reacher *et al.* 2004; Paranjothy *et al.* 2011; Fontalba-Navas *et al.* 2017; Milojevic *et al.* 2011). Inundation of property, damage to possessions and forced evacuations due to flooding can all be classified as direct disruptions. A key finding regarding direct disruptions has been their significant association with post-traumatic stress disorder (PTSD) (Waite *et al.* 2017; Munro *et al.* 2017; Jermacane *et al.* 2018; Zhong *et al.* 2018; Alderman, Turner & Tong 2013; Paranjothy *et al.* 2011; Matthews *et al.* 2019)

Conversely, there is a dearth of published research investigating the associations between mental health morbidity and *indirect* disruptions such as losing access to health care, food or place of employment for people who have not had their property inundated with flood water. Two of the only studies of indirect disruptions following flooding events come from England. These studies identified adverse effects on mental health, especially PTSD (Paranjothy *et al.* 2011, Waite *et al.* 2017). To date, there has been no research on the mental health effects of indirect disruptions published in the Australian context.

Objectives

The objectives of this research:

- Examine the associations between an experience of indirect disruption and direct disruption with mental health outcomes after a flood.
- Examine which types of indirect disruption due to flooding are most strongly associated with adverse mental health outcomes.

Study design

From September to November 2017, six months after ex-Tropical Cyclone Debbie caused extensive flooding

in northern New South Wales, a cross-sectional survey was implemented targeting people who had been living in six local government areas of Ballina Shire, Tweed Shire, Richmond Valley, Kyogle, Byron Shire and Lismore City at the time of the flood. These areas had an estimated population of 247,000 (Australian Bureau of Statistics 2017). Community members aged 16 years and older were recruited using a 'snowball' sampling method. This method incorporated social and organisational networks of local government authorities, business groups and community organisations and was supplemented by an extensive local advertising campaign using print, broadcast and social media. This included a leaflet drop in the two largest centres of population flooded; Lismore and Murwillumbah. All residents were encouraged to participate whether or not they felt the flood had affected them. The survey was available in online and paper formats. Potential respondents were advised that completion of the questionnaire would signify consent to participate in the study. A detailed description of the study design was published by Longman and colleagues (2019).

The study was approved by the University of Sydney Human Research Ethics Committee (reference-2017/589) and the Aboriginal Health and Medical Research Council Human Research Ethics Committee (reference-1294/17).

Measures of disruption

The survey contained questions relating to the degree of flooding, disruption, socio-demographic characteristics and the mental health and wellbeing of the respondents. Using the responses to the disruption questions, participants were categorised into three independent groups: directly disrupted, indirectly disrupted and non-disrupted (Waite *et al.* 2017).

Directly disrupted respondents: Those who reported flooding or damage to any area of their home or income-generating property (e.g. business or farm). These respondents had answered yes to at least one of the following questions:

- *Were non-livable areas of your home damaged or flooded (e.g. garage, garden shed)?*
- *Was at least one livable room in your home damaged or flooded (e.g. bedroom, living room, kitchen, bathroom)?*
- *If you own a business, was it damaged or flooded (e.g. if you own a shop, farm, warehouse)?*

Indirectly disrupted respondents: Those who were not flooded (i.e. answered 'No' to all the previous questions) but who answered yes to any of the following:

- *Your access to health and social care was disrupted.*
- *You had difficulty getting the food supplies you needed.*
- *You were temporarily isolated as surrounding roads were cut.*
- *You were unable to travel to your place of education (e.g. school, university, TAFE).*
- *There were interruptions to your household utilities (e.g. electricity, gas, drainage, septic).*
- *Your Wi-Fi/internet stopped working.*
- *You were unable to travel to your place of employment.*

Non-disrupted respondents: Those who reported none of the specified disruptions.

Measures to assess mental health

Previous studies of natural disasters including flooding have indicated that PTSD, depression, anxiety and increased suicide risk are common sequelae (Alderman, Turner & Tong 2013; Paranjothy *et al.* 2011; Waite *et al.* 2017; Zhong *et al.* 2018; Tang *et al.* 2018; Lowe *et al.* 2019). Therefore, in this study, mental health status was assessed using brief versions of validated screening tools of:

- the Patient Health Questionnaire (PHQ-2) for depression (Kroenke, Spitzer & Williams 2003)
- the Generalised Anxiety Disorder scale (GAD-2) (Kroenke *et al.* 2007)
- the Post Traumatic Stress Disorder Checklist (PCL-6) (Lang & Stein 2005, Fernandez *et al.* 2015).

The PHQ-2, GAD-2 and PCL-6 were selected to keep the outcomes of this study in-line with the only other study previously published investigating the comparison between disruption type due to flooding and adverse mental health outcomes (Waite *et al.* 2017).

Cut-points for probable diagnosis were ≥ 3 for the PHQ-2 and GAD-2 and ≥ 14 for the PCL-6 (Lang & Stein 2005; Kroenke, Spitzer & Williams 2003; Kroenke *et al.* 2007). To relate responses to the PCL-6 to the flood, the checklist was introduced as a list of complaints that people express after extreme rain and flooding. Additional mental health measures included an indicator of suicidal ideation from the Screening Tool for Assessing Risk of Suicide (Hawgood & DeLeo 2017) and an indicator of continuing distress six months after the flood (Clemens *et al.* 2013).

Socio-demographic measures

Socio-demographic data included age, gender, Indigenous status, relationship status, education level, employment status and government income support status. Only respondents with complete socio-demographic data were included in the analysis.

Analysis by disruption category

The analysis was conducted in two stages. First, binary logistic regression models were constructed to calculate the odds of experiencing each of the five types of mental health outcomes: continuing distress, suicidal ideation, probable depression, anxiety and PTSD. Respondents who did not complete a particular health outcome measure were excluded from analysis for that outcome. The dependent variables were the category of disruption, with non-disruption as the reference group. The models were adjusted for all measured socio-demographic characteristics. Sensitivity analyses were conducted to assess the level of bias introduced by including these characteristics.

Analysis by type of indirect disruption

In the second stage of the analysis, five multivariate logistic regression models were constructed that considered the

association between each type of indirect disruption with each of the five mental health conditions as an outcome.

It was reasoned that a substantial portion of participants who experienced an indirect disruption was likely to also have experienced direct disruption. Therefore, only analysing the participants who experienced an indirect disruption without direct disruption would have resulted in a markedly reduced sample from which conclusions could be drawn and could introduce risk of bias in the results. For this reason, every participant who reported an indirect disruption was included and participants were not grouped by disruption category. To account for potential confounding caused by experiencing both direct and indirect disruption, the regression models included a binary variable that indicated any experience of direct disruption by the participants. Again, each model was adjusted for socio-demographic characteristics.

The interest was in identifying significant associations and important confounders. As such, purposeful selection was employed to construct the multivariate logistic regression models. Consistent with the purposeful selection method, other indirect disruptors were retained in the model if they demonstrated a *p*-value of less than 0.15 or if they demonstrated significant confounding effects ($\Delta\beta > 20\%$) (Bursac *et al.* 2008). As there were multiple analyses investigating the mental health outcomes in both sets of analyses, the α for significance testing was set conservatively at 0.01. Every regression model produced in this study was tested for effect modifications ($\alpha=0.01$). Stata 15 (Stata/SE 15.1 for Windows) was used for all statistical analyses.

Results

Respondent characteristics

In total, 2530 people responded to the survey and 350 (14 per cent) of the responses were missing socio-demographic data. Therefore, the analysis conducted using socio-demographic data was performed using a sample of 2180 participants. Negligible dissimilarities in parameter estimates and patterns of results were found between the full dataset and the dataset absent of missing socio-demographic records.

Mental health outcomes by disruption category

Of these 2180 respondents, 105 could not be classified into disruption categories due to incomplete survey responses, 242 respondents were classified as non-disrupted, 605 were classified as indirectly disrupted and 1228 were classified as directly disrupted. In total, 2075 respondents were included in this part of the analysis. Most of the respondents were over 45 years of age, were female, in a relationship and employed (Table 1, Appendix 1).

Among those who were classified as directly disrupted, between 10 per cent and 33 per cent demonstrated evidence of mental health distress in the outcomes measured. By comparison, among those who were classified as non-disrupted, between 2 and 8 per cent reported mental health distress (Table 2).

Accordingly, the differences in proportions demonstrating evidence of mental health distress between the two groups ranged from 7 per cent (suicidal ideation) to 25 per cent (still distressed).

The differences in proportions demonstrating evidence of mental health distress between those classified as indirectly disrupted and non-disrupted was less stark, ranging from 1 per cent (suicidal ideation) to 7 per cent (probable anxiety).

When factoring in potential confounders in the logistic regression model, the greatest effect on the odds of probable PTSD was found in respondents who experienced direct disruption (OR: 14.4; 95 per cent, CI 5.9–35.3) (Table 3). After adjusting for socio-demographic factors, probable PTSD remained strongly associated with direct disruption (OR:13.5; 95 per cent, CI: 5.5–33.4). Indeed, the odds of experiencing every mental health outcome remained significantly elevated in response to direct disruption after adjusting for socio-demographic factors (Table 3).

Respondents categorised as indirectly disrupted were significantly more likely to experience probable PTSD, probable anxiety or still feel distressed than those who were categorised as non-disrupted. After adjusting for socio-demographic factors, only the odds of probable PTSD remained significantly elevated (OR: 3.52, 95 per cent, CI: 1.36–9.15) (Table 3).

Mental health outcome by indirect disruption type

The relationships between each mental health outcome and each of the indirect disruption types were also examined using multivariate logistic regressions (*n*=2180). The other indirect disruptions were added to the models according to the purposeful selection method and were also adjusted for any experience of direct disruption and socio-demographic factors.

Loss of access to social or health care was shown to significantly increase the odds of every outcome except probable depression. No individual disruption type significantly influenced the odds of having probable depression (Table 5). On the outcome of probable anxiety, employment status significantly modified the size of the effect of a loss of access to health and social care. Among those participants who experienced a loss of access to health and social care, unemployed participants demonstrated greater odds of probable anxiety than those who were employed (OR: 2.67; 95 per cent, CI: 1.64, 4.35 vs OR: 1.05; 95 per cent, CI: 0.67, 1.64, respectively). A loss of utilities was strongly associated with every mental health outcome although this effect was only statistically significant for probable PTSD (OR: 1.9, 95 per cent, CI: 1.41–2.56) (Table 5).

Discussion

The strong link between disruption after a flood event and PTSD has been clearly elucidated in recent literature (Fontalba-Navas *et al.* 2017, Zhong *et al.* 2018, Dai *et al.* 2017, Waite *et al.* 2017, Paranjothy *et al.* 2011, Fernandez *et al.* 2015, Matthews *et al.* 2019). Consistent with Waite and co-authors (2017), these results

Table 1: Socio-demographic characteristics of respondents by disruption category.

Demographic		Disruption Category			
		No disruption	Indirectly	Directly	Total
Age (years)	16 to 25	8 (3%)	37 (6%)	65 (5%)	110 (5%)
	24 to 45	51 (21%)	173 (29%)	290 (24%)	514 (25%)
	45 to 65	115 (48%)	296 (49%)	667 (54%)	1078 (52%)
	65 and older	68 (28%)	99 (16%)	206 (17%)	373 (18%)
	Total	242 (100%)	605 (100%)	1228 (100%)	2075 (100%)
Gender	Female	163 (67%)	430 (71%)	834 (68%)	1427 (69%)
	Male	79 (33%)	175 (29%)	394 (32%)	648 (31%)
	Total	242 (100%)	605 (100%)	1228 (100%)	2075 (100%)
Aboriginal or Torres Strait Islander peoples	Yes	9 (4%)	14 (2%)	53 (4%)	76 (4%)
	No	233 (96%)	591 (98%)	1175 (96%)	1999 (96%)
	Total	242 (100%)	605 (100%)	1228 (100%)	2075 (100%)
Relationship status	In a relationship	162 (67%)	414 (68%)	820 (67%)	1396 (67%)
	Single	80 (33%)	191 (32%)	408 (33%)	679 (33%)
	Total	242 (100%)	605 (100%)	1228 (100%)	2075 (100%)
Education attained	University	119 (49%)	304 (50%)	483 (39%)	906 (44%)
	Other	123 (51%)	301 (50%)	745 (61%)	1169 (56%)
	Total	242 (100%)	605 (100%)	1228 (100%)	2075 (100%)
Employment status	Employed	143 (59%)	450 (74%)	835 (68%)	1428 (69%)
	Other	99 (41%)	155 (26%)	393 (32%)	648 (31%)
	Total	242 (100%)	605 (100%)	1228 (100%)	2075 (100%)
Income support	None	165 (68%)	451 (75%)	800 (65%)	1416 (68%)
	Support	77 (32%)	154 (25%)	428 (35%)	659 (32%)
	Total	242 (100%)	605 (100%)	1228 (100%)	2075 (100%)

Table 2: Respondent mental health outcome by disruption category.

Outcome	Overall cohort	Disruption group		
		Non-disrupted	Indirectly	Directly
Still distressed	478/2050 (23%)	20/242 (8%)	58/597 (10%)	400/1211 (33%)
Probable PTSD	327/2044 (16%)	5/232 (2%)	40/599 (7%)	282/1213 (23%)
Depression	326/2026 (16%)	13/235 (6%)	57/590 (10%)	256/1201 (21%)
Probable anxiety	335/2021 (17%)	9/232 (4%)	62/590 (11%)	264/1199 (22%)
Suicidal Ideation	156/2056 (8%)	8/240 (3%)	23/601 (4%)	125/1215 (10%)
Total*	2075	242	605	1228

* Totals differ from the overall sample size due to missing responses for outcome and exposure reporting.

Table 3: Crude and adjusted odds ratios (OR) for mental health problems by disruption category.

Outcome	Disruption group	n	Crude OR	Crude p-value	n	Adjusted OR [#]	Adjusted p-value [#]
Still distressed	Non-disrupted	260	1 [†]	-	242	1 [†]	-
	Indirectly	654	1.21 (0.72–2.05)	<0.001	597	0.95 (0.47–1.89)	0.878
	Directly	1374	5.77 (3.6–9.23)	<0.001	1211	3.31 (1.79–6.12)	<0.001
Probable PTSD	Non-disrupted	246	1 [†]	-	232	1 [†]	-
	Indirectly	646	3.52 (1.38–8.99)	0.008	599	3.52 (1.36–9.15)	0.01
	Directly	1341	14.43 (5.9–35.31)	<0.001	1213	13.48 (5.45–33.35)	<0.001
Probable depression	Non-disrupted	249	1 [†]	-	235	1 [†]	-
	Indirectly	639	1.88 (1.01–3.49)	0.045	590	1.9 (1–3.62)	0.05
	Directly	1331	4.99 (2.81–8.86)	<0.001	1201	4.26 (2.35–7.73)	<0.001
Probable anxiety	Non-disrupted	247	1 [†]	-	232	1 [†]	-
	Indirectly	639	2.56 (1.33–4.92)	0.005	590	1.55 (0.7–3.44)	0.279
	Directly	1325	6.09 (3.28–11.3)	<0.001	1199	3.64 (1.74–7.62)	0.001
Suicidal ideation	Not Disrupted	256	1 [†]	-	240	1 [†]	-
	Indirectly	648	1.24 (0.55–2.8)	0.597	601	1.02 (0.44–2.35)	0.961
	Directly	1340	3.47 (1.68–7.18)	0.001	1215	2.86 (1.36–5.99)	0.005

Adjusted for age, gender, Indigenous status, receiving income support, education and relationship status.

† Reference group.

showed a significant association between direct disruption and probable PTSD. By comparison, significant association has also been demonstrated between indirect disruption and probable PTSD, although the strength of association is comparatively weaker. This apparent dose-response relationship offers evidence for the causative relationship between the level of disruption due to flooding and the outcome of probable PTSD.

Direct disruption also demonstrated significant associations with the other four mental health outcomes when compared with non-disruption, namely: still distressed, probable depression, probable anxiety and suicidal ideation. By contrast, no significant associations were found between indirect disruption and these four mental health outcomes.

This is one of the few studies that has sought to investigate the associations between mental health and indirect disruption due to a flooding event. Consistent with previous research, indirect disruption was found to be significantly associated with an increased risk of probable PTSD in comparison with individuals classified as non-disrupted (Waite *et al.* 2017, Paranjothy *et al.* 2011).

To date, there has been little discussion in the literature about mechanisms that might account for this increased risk of probable PTSD in cases where there has been disruption

experienced, albeit with no direct damage to people’s homes or businesses. Some indications of possible mechanisms might be derived from research on the impact of near-miss experiences and PTSD diagnostic criteria.

Recent literature on ‘near-miss experiences’ following traumatic events suggests that people who have had near-miss events tend to experience more intrusive thoughts about what might have been and are more likely to think about the actual misfortune of others, which may reinforce intrusions and raise the likelihood of post-traumatic stress symptoms (Poulin & Silver 2019). It might be inferred that those who experienced indirect disruption had a ‘near-miss experience’ and may have been more sensitive to what might have been and therefore more prone to post-traumatic stress than those who were classified as non-disrupted.

Except for probable depression, these results demonstrate a strong association between losing access to health and social care and every mental health outcome investigated. A similar association was reported by Waite and colleagues (2017), although their results were not mutually adjusted for other disruption types. Interruption to household utilities was also shown to significantly increase the odds of having probable PTSD, consistent with similar findings reported related to the loss of electricity after a flooding event in Hat Yai, Thailand (Assanangkornchai, Tangboonngam & Edwards 2004).

Table 4: Multivariate logistic regression models for mental health outcomes and individual indirect disruption types.

Mental health outcome	Contributing variable	Adjusted odds ratio (95% CI)	P-value
Still distressed	Loss of access to social or health care		
	No loss of access	1	-
	Loss of access**	1.86 (1.38–2.49)	<0.001
	Loss of utilities		
	No loss of access	1	-
	Loss of access	1.34 (1.02–1.75)	0.034
	Loss of access to internet		
	No loss of internet	1	-
Loss of internet	1.4 (1.08–1.82)	0.012	
Probable PTSD	Loss of access to social or health care		
	No loss of Access	1	-
	Loss of access**	1.93 (1.38–2.7)	<0.001
	Loss of utilities		
	No loss of utilities	1	-
	Loss of utilities**	1.9 (1.41–2.56)	<0.001
	Difficulty accessing food		
	No difficulty	1	-
Difficulty	1.31 (0.94–1.83)	0.105	
Probable depression	Loss of access to social or health care		
	No loss of access	1	-
	Loss of access	1.51 (1.44–2.82)	0.016
	Loss of utilities		
	No loss of utilities	1	-
	Loss of utilities	1.38 (1.02–1.85)	0.035
Probable anxiety [^]	Loss of access to social or health care		
	Not employed/No loss of access	1	-
	Not employed/Loss of access**	2.67 (1.64–4.35)	<0.001
	Employed/No loss of access	1	-
	Employed/Loss of access	1.05 (0.67–1.64)	0.838
	Loss of utilities		
	No loss of access	1	-
	Loss of access	1.31 (0.98–1.74)	0.07
Suicide ideation	Loss of access to social or health care		
	No loss of access	1	-
	Loss of access*	1.74 (1.14–2.66)	0.01
	Loss of utilities		
	No loss of access	1	-
Loss of access	1.43 (0.96–2.13)	0.079	

Each model is adjusted for age, gender, Indigenous status, receiving income support, education, relationship status and an experience of direct disruption; remaining covariates retained and presented as per the purposeful selection method.

[^]Denotes significant effect modification identified ($\alpha=0.01$)

* Denotes p -value ≤ 0.01 .

** Denotes p -value ≤ 0.001 .

People who lost access to health and social care as well as being unemployed, were more likely to have an outcome of probable anxiety than those who were employed. It may be that people experiencing unemployment are more likely to need access to health and social care than people who are employed. It may also be that unemployment as well as a loss of this access may have a cumulative effect on anxiety.

There were increased odds of reporting probable PTSD for participants who were classified as indirectly disrupted and for those who specifically reported either a disruption of access to social and health care or a disruption to utilities. Therefore, it is possible that the association between an experience of indirect disruption and the outcome of probable PTSD is largely explained by losing access to social or health care and/or a disruption to utilities.

Given the predicted intensification of the effects of climate change (Climate Council 2017), there is likely to be an increasing number of people who are disrupted by flooding events. This study offers evidence that after a flooding event those in need of mental health support will include people who have been directly disrupted and also those who have lost access to social and health care as well as those who have experienced a disruption to utilities provision.

Treatment of mental health problems, including PTSD, following disasters requires specific training for those working in the mental health field (Foa, Gillihan & Bryant 2013). Furthermore, it is recognised that different approaches delivered by professionals may be needed at different stages post disaster (Forbes, O'Donnell & Bryant 2017). Given the findings of this study, it may be prudent to provide the health care workforce with access to appropriate-level training (e.g. mental health first-aid for the general health and community sector, PTSD training for specialists) in preparation for future acute events. Part of this training could involve how to identify people most in need of mental health care, including those who have been indirectly disrupted. Also, given that those who require mental health support after a flood event may need assistance more than six months after a flooding event (Zhong *et al.* 2018), planning is needed to provide support in the longer, not just immediate, term.

Limitations

The self-selection recruitment method means the respondent population is not representative of the population of the flood-affected communities of northern New South Wales. In this study, men, those with less education and those under 25 years of age were under-represented, which is consistent with previous post-disaster postal survey respondent characteristics (Grievink *et al.* 2006). However, this study did not aim to establish prevalence of mental health outcomes after flooding, but rather to examine the relationships between levels and types of disruption and mental health status. Further, to ascertain the relationships of interest, a study's population does not necessarily have to be representative of the general population from which it was derived (Willett *et al.* 2007, Banks *et al.* 2008).

Conclusions

The responses to the community survey conducted in northern New South Wales have contributed to knowledge around mental health effects after a significant flooding event (Matthews *et al.* 2019). The findings are consistent with, and extend, what has been presented in recent literature. It is proposed that people experiencing disruptions to social and health care services or utilities after a flood might be targeted for mental health support.

Further research in this area might investigate causal mechanisms behind indirect disruptions and their associated mental health outcomes. Also, planned follow-up surveys involving respondents who have indicated an interest in future participation may shed light on the longer-term mental health consequences of flooding events.

References

- Alderman K, Turner LR & Tong S 2012, *Assessment of the health impacts of the 2011 summer floods in Brisbane*. NCCARF National Adaptation Conference. Melbourne.
- Alderman K, Turner LR & Tong S 2013, *Assessment of the health impacts of the 2011 summer floods in Brisbane*. *Disaster Medicine and Public Health Preparedness*, 7, pp.380–386.
- Assanangkornchai S, Tangboonngam S-N & Edwards JG 2004, *The flooding of Hat Yai: predictors of adverse emotional responses to a natural disaster*. *Stress and Health*, vol. 20, pp.81–89.
- Australian Bureau of Statistics 2017, *Estimated resident population (ABS consultancy), Statistical area level 1 (SA1), Australia, 2011 to 2016p*. Queensland Government Statisticians Office. At: www.qgso.qld.gov.au/products/tables/erp-sa1-aus-consult/index.php?region=brisbane [January 2018].
- Banks E, Redman S, Jorm L, Armstrong B, Bauman A, Beard J, Beral V, Byles J, Corbett S, Cumming R, Harris M, Sitas F, Smith W, Taylor L, Wutzke S & Lujic S 2008, *Cohort profile: the 45 and up study*. *International Journal of Epidemiol*, vol. 37, pp.941–947. doi 10.1093/ije/dym184
- Bursac Z, Gauss CH, Williams DK & Hosmer DW 2008, *Purposeful selection of variables in logistic regression*. *Source Code for Biology and Medicine*, vol. 3, p.17. At: <https://scfbm.biomedcentral.com/articles/10.1186/1751-0473-3-17>.
- Clemens SL, Berry HL, McDermott BM & Harper CM 2013, *Summer of sorrow: measuring exposure to and impacts of trauma after Queensland's natural disasters of 2010–2011*. *Medical Journal of Australia*, vol. 199, pp.552–555.
- Climate Council 2017, *Intense Rainfall and Flooding: the influence of climate change Sydney, Australia*.
- Dai W, Kaminga AC, Tan H, Wang J, Lai Z, Wu X & Liu A 2017, *Long-term psychological outcomes of flood survivors of hard-hit areas of the 1998 Dongting Lake flood in China: Prevalence and risk factors*. *PLoS One*, vol. 12, e0171557.
- Fernandez A, Black J, Jones M, Wilson L, Salvador-Carulla L, Astell-Burt T & Black D 2015, *Flooding and Mental Health: A Systematic Mapping Review*. *PLoS ONE*, vol. 10, e0119929.
- Foa EB, Gillihan SJ & Bryant RA 2013, *Challenges and Successes in Dissemination of Evidence-Based Treatments for Posttraumatic Stress: Lessons Learned From Prolonged Exposure Therapy for PTSD*. *Psychological Science in the Public Interest*, vol. 14, pp.65–111.

- Fontalba-Navas A, Lucas-Borja ME, Gil-Aguilar V, Arrebola JP, Pena-Andreu JM & Perez J 2017, *Incidence and risk factors for post-traumatic stress disorder in a population affected by a severe flood. Public Health*, vol. 144, pp.96–102.
- Forbes D, O'Donnell M & Bryant RA 2017, *Psychosocial recovery following community disasters: An international collaboration. Australian New Zealand Journal of Psychiatry*, vol. 51, pp.660–662.
- Grievink L, van der Velden PG, Yzermans CJ, Roorda J & Stellato RK 2006, *The importance of estimating selection bias on prevalence estimates shortly after a disaster. Ann Epidemiol*, vol. 16, pp.782–788.
- Hawgood J & Deleo D 2017, *STARS: Screening Tool for Assessing Risk of Suicide. Brisbane: Australian Institute for Suicide Research and Prevention, Griffith University.*
- Jermacane D, Waite TD, Beck CR, Bone A, Amlôt R, Reacher M, Kovats S, Armstrong B, Leonardi G, James Rubin G & Oliver I 2018, *The English National Cohort Study of Flooding and Health: the change in the prevalence of psychological morbidity at year two. BMC Public Health*, vol. 18, p.330.
- Kroenke K, Spitzer R & Williams J 2003, *The Patient Health Questionnaire-2: Validity of a Two-item Depression Screener. Medical Care*, vol. 41, pp.1284–1292.
- Kroenke K, Spitzer R, Williams J, Monahan P & Lowe B 2007, *Anxiety Disorders in Primary Care: Prevalence, Impairment, Comorbidity, and Detection. Annals of Internal Medicine*, vol. 146, pp.317–325.
- Lang AJ & Stein MB 2005, *An abbreviated PTSD checklist for use as a screening instrument in primary care. Behaviour Research and Therapy*, vol. 43, pp.585–594.
- Longman JM, Bennett-Levy J, Matthews V, Berry HL, Passey ME, Rolfe M, Morgan GG, Braddon M & Bailie R 2019, *Rationale and methods for a cross-sectional study of mental health and wellbeing following river flooding in rural Australia, using a community-academic partnership approach. BMC Public Health*, vol. 19, p.1255.
- Lowe SR, Bonumwezi JL, Valdespino-Hayden Z & Galea S 2019, *Posttraumatic Stress and Depression in the Aftermath of Environmental Disasters: A Review of Quantitative Studies Published in 2018. Current Environmental Health Reports*.
- Matthews V, Longman JM, Berry HL, Passey ME, Bennett-Levy J, Morgan GG, Pit S, Rolfe M, Bailie R 2019, *Differential mental health impact six months after extensive river flooding in rural Australia: a cross-sectional analysis through an equity lens. Frontiers of Public Health*, vol. 7, p.367.
- Milojevic A, Armstrong B, Kovats S, Butler B, Hayes E, Leonardi G, Murray V & Wilkinson P 2011, *Long-term effects of flooding on mortality in England and Wales, 1994–2005: controlled interrupted time-series analysis. Environmental Health*, vol. 10, p.11.
- Munro A, Kovats RS, Rubin GJ, Waite TD, Bone A, Armstrong B & English National Study of Flooding and Health Study Group 2017, *Effect of evacuation and displacement on the association between flooding and mental health outcomes: a cross-sectional analysis of UK survey data. Lancet Planet Health*.
- Paranjothy S, Gallacher J, Amlôt R, Rubin GJ, Page L, Baxter T, Wight J, Kirrage D, MCNaught R & Sr P 2011, *Psychosocial impact of the summer 2007 floods in England. BMC Public Health*, vol. 11, p.145.
- Poulin MJ & Silver RC 2019, *What Might Have Been: Near Miss Experiences and Adjustment to a Terrorist Attack. Social Psychological and Personality Science*. doi: 10.1177/1948550619829064
- Reacher M, McKenzie K, Lane C, Nichols T, Kedge I, Iversen A, Hepple P, Walter T, Laxton C & Simpson J 2004, *Health impacts of flooding in Lewes: a comparison of reported gastrointestinal and other illness and mental health in flooded and non-flooded households. Communicable Disease and Public Health*, vol. 7, pp.39–46.
- Tang W, Xu D, Li B, Lu Y & Xu J 2018, *The relationship between the frequency of suicidal ideation and sleep disturbance factors among adolescent earthquake victims in China. General Hospital Psychiatry*, vol. 55, pp.90–97.
- Waite TD, Chaintarli K, Beck CR, Bone A, Amlôt R, Kovats S, Reacher M, Armstrong B, Leonardi G, Rubin GJ & Oliver I 2017, *The English national cohort study of flooding and health: cross-sectional analysis of mental health outcomes at year one. BMC Public Health*, vol. 17, p.129.
- Willett WC, Blot WJ, Colditz GA, Folsom AR, Henderson BE & Stampfer MJ 2007, *Merging and emerging cohorts: not worth the wait. Nature*, vol. 445, pp.257–258.
- Zhong S, Yang L, Toloo S, Wang Z, Tong S, Sun X, Crompton D, Fitzgerald G & Huang C 2018, *The long-term physical and psychological health impacts of flooding: A systematic mapping. Science of the Total Environment*, vol. 626, pp.165–194.

About the authors

Jonathan King is an epidemiologist at the Kirby Institute, University of New South Wales. His interests are in mental health and preventable hospitalisation.

Dr Jo Longman is a research fellow at the University of Sydney, University Centre for Rural Health. Her interests include the mental health effects of weather-related and acute disasters.

Dr Veronica Matthews is a researcher at the University of Sydney, University Centre for Rural Health. Her interests are in community-based research methods and improving health care systems in Aboriginal and Torres Strait Islander communities.

Professor James Bennett-Levy is a mental health researcher at the University of Sydney, University Centre for Rural Health. His interests are in enhancing the social and emotional wellbeing of Aboriginal and Torres Strait Islander peoples and people living in regional and rural Australia.

Professor Ross Bailie is the Director of the University of Sydney, University Centre for Rural Health. He has qualifications in medicine, general practice, maternal and child health and public health and has worked in South Africa, New Zealand and Australia.

Steve Carrigg is a registered nurse and has worked in acute mental health services for 30 years, primarily in rural and remote Australia. He is the coordinator of the Rural Adversity Mental Health Program for the Northern New South Wales Local Health District. He is involved in the coordination of the mental health response during disasters and ongoing recovery.

Associate Professor Megan Passey is the Director of Research at the University of Sydney, University Centre for Rural Health. Her interests include ways to address complex issues in rural and Indigenous health.

Appendix i. Number and proportions of respondents reporting mental health outcomes for each indirect disruption type.

	Loss of access to food		Loss of access to social/health care		Isolated due to flooding		Couldn't get to education		Disrupted utilities		Loss of access to internet		Couldn't get to work	
	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)
Still	132 (30)	354 (21)	135 (41)	351 (19)	328 (27)	100 (15)	94 (30)	334 (22)	244 (34)	184 (16)	188 (32)	240 (19)	191 (24)	237 (22)
distressed	No	302 (70)	1363 (79)	1468 (81)	870 (73)	560 (85)	217 (70)	1213 (78)	476 (66)	954 (84)	407 (68)	1023 (81)	612 (76)	818 (78)
Total	434 (100)	1717 (100)	332 (100)	1819 (100)	1198 (100)	660 (100)	311 (100)	1547 (100)	720 (100)	1138 (100)	595 (100)	1263 (100)	803 (100)	1055 (100)
Probable	107 (25)	225 (13)	115 (35)	217 (12)	231 (19)	62 (9)	70 (22)	223 (14)	189 (26)	104 (9)	130 (22)	163 (13)	128 (16)	165 (16)
PTSD	No	324 (75)	1458 (87)	1565 (88)	971 (81)	591 (91)	242 (78)	1320 (86)	536 (74)	1026 (91)	461 (78)	1101 (87)	674 (84)	888 (84)
Total	431 (100)	1683 (100)	332 (100)	1782 (100)	1202 (100)	653 (100)	312 (100)	1543 (100)	725 (100)	1130 (100)	591 (100)	1264 (100)	802 (100)	1053 (100)
Probable	97 (23)	237 (14)	95 (29)	239 (14)	204 (17)	82 (13)	65 (21)	221 (14)	158 (22)	128 (11)	109 (19)	177 (14)	105 (13)	181 (17)
depression	No	328 (77)	1435 (86)	1530 (86)	982 (83)	569 (87)	243 (79)	1308 (86)	562 (78)	989 (89)	477 (81)	1074 (86)	685 (87)	866 (83)
Total	425 (100)	1672 (100)	328 (100)	1769 (100)	1186 (100)	651 (100)	308 (100)	1529 (100)	720 (100)	1117 (100)	586 (100)	1251 (100)	790 (100)	1047 (100)
Probable	104 (24)	238 (14)	102 (31)	240 (14)	219 (18)	81 (13)	75 (24)	225 (15)	164 (23)	136 (12)	115 (20)	185 (15)	123 (16)	177 (17)
anxiety	No	321 (76)	1432 (86)	1528 (86)	966 (82)	566 (87)	234 (76)	1298 (85)	551 (77)	981 (88)	471 (80)	1061 (85)	670 (84)	862 (83)
Total	425 (100)	1670 (100)	327 (100)	1768 (100)	1185 (100)	647 (100)	309 (100)	1523 (100)	715 (100)	1117 (100)	586 (100)	1246 (100)	793 (100)	1039 (100)
Suicidal	Yes	52 (12)	106 (6)	108 (6)	105 (9)	32 (5)	37 (12)	100 (6)	79 (11)	58 (5)	50 (8)	87 (7)	61 (8)	76 (7)
ideation	No	377 (88)	1595 (94)	1691 (94)	1092 (91)	635 (95)	276 (88)	1451 (94)	645 (89)	1082 (95)	546 (92)	1181 (93)	745 (92)	982 (93)
Total	429 (100)	1701 (100)	331 (100)	1799 (100)	1197 (100)	667 (100)	313 (100)	1551 (100)	724 (100)	1140 (100)	596 (100)	1268 (100)	806 (100)	1058 (100)
All	434	1717	332	1819	1210	671	314	1567	730	1151	601	1280	811	1070